

CURRENT CONCEPTS REVIEW**Posterior Cruciate Ligament: Current Concepts Review**

Santiago Pache, MD; Zachary S. Aman, BA; Mitchell Kennedy, BA; Gilberto Yoshinobu Nakama MD; Gilbert Moatshe, MD; Connor Ziegler, MD; Robert F LaPrade, PhD

Research performed at the Steadman Philippon Research Institute, Vail, Colorado, USA

Received: 30 August 2017

Accepted: 29 November 2017

Abstract

The posterior cruciate ligament (PCL) is the largest and strongest ligament in the human knee, and the primary posterior stabilizer. Recent anatomy and biomechanical studies have provided an improved understanding of PCL function. PCL injuries are typically combined with other ligamentous, meniscal and chondral injuries. Stress radiography has become an important and validated objective measure in surgical decision making and post-operative assessment. Isolated grade I or II PCL injuries can usually be treated non-operatively. However, when acute grade III PCL ruptures occur together with other ligamentous injury and/or repairable meniscal body/root tears, surgery is indicated. Anatomic single-bundle PCL reconstruction (SB-PCLR) typically restores the larger anterolateral bundle (ALB) and represents the most commonly performed procedure. Unfortunately, residual posterior and rotational tibial instability after SB-PCLR has led to the development of an anatomic double-bundle (DB) PCLR to restore the native PCL footprint and co-dominant behavior of the anterolateral and posteromedial bundles and re-establish normal knee kinematics. The purpose of this article is to review the pertinent details regarding PCL anatomy, biomechanics, injury diagnosis and treatment options, with a focus on arthroscopically assisted DB-PCLR.

Level of evidence: IV

Keywords: Double bundle posterior cruciate ligament reconstruction, Posterior cruciate ligament, Posterior knee laxity, Stress radiographs

Introduction

Posterior cruciate ligament (PCL) tears comprise 3% of outpatient knee injuries and 38% of acute traumatic knee hemarthroses (1). These injuries rarely occur in isolation, and up to 95% of PCL tears occur in combination with other ligament tears. With more people participating in sporting activities, these injuries will potentially increase in the future. PCL tears are increasingly being recognized as source of morbidity and reduced function because of persistent instability, pain, impaired function and development of degenerative joint disease (2, 3).

In recent years, a better understanding of the anatomy and biomechanics of the PCL has emerged, leading to improved surgical techniques and rehabilitation protocols for the treatment of PCL tears. However, controversy still exists regarding the decisions for non-

operative versus operative treatment, and the optimal surgical technique. The heterogeneity of these injuries can make it difficult to compare studies in the literature. Furthermore, studies with long-term follow-up are still lacking in the literature. The purpose of this review was to report on the current concepts of PCL tears including the anatomy, biomechanics, diagnosis, treatment options, rehabilitation, and outcomes reported in the literature.

Anatomy

The PCL is the largest and strongest intraarticular ligament of the knee joint, comprising of 2 functional bundles: the larger anterolateral bundle (ALB) and the smaller posteromedial bundle (PMB) (4). The size of the femoral attachment of the ALB is nearly twice the size

Corresponding Author: Robert F. LaPrade, Steadman Philippon Research Institute, The Steadman Clinic, Vail, Colorado, USA
Email: drlaprade@srpivail.org



THE ONLINE VERSION OF THIS ARTICLE
[ABJS.MUMS.AC.IR](http://abjs.mums.ac.ir)




Figure 1. (A) Anterior and (B) posterior views of the native posterior cruciate ligament (PCL). Emphasized are the femoral and tibial attachments of the anterolateral bundle (ALB) and posteromedial bundle (PMB) of the PCL and the osseous landmarks: the trochlear point, the medial arch point, the bundle ridge, and the champagne-glass drop-off. ACL, anterior cruciate ligament; aMFL, anterior meniscofemoral ligament (ligament of Humphrey); FCL, fibular collateral ligament; PFL, popliteofibular ligament; pMFL, posterior meniscofemoral ligament (ligament of Wrisberg); POL, posterior oblique ligament (*Reproduced with permission from Kennedy NJ, Wijdicks CA, Goldsmith MT, et al. Kinematic analysis of the posterior cruciate ligament, part 1: the individual and collective function of the anterolateral and posteromedial bundles. Am J Sports Med. 2013;41(12):2828-2838.*).

of its tibial attachment and has been reported to range from 112 to 118 mm² (5–7). The center of the femoral ALB footprint is located 7.4 mm from the trochlear point, 11.0 mm from the medial arch point, and 7.9 mm from the distal articular cartilage. Furthermore, ALB tibial attachment center is located 6.1 mm posterior to the shiny white fibers of the posterior medial meniscus root, 4.9 mm from the bundle ridge (which separates both bundles), and 10.7 mm from the champagne glass drop-off of the posterior tibia (5).

The area of the PMB femoral attachment is between 60 mm² and 90 mm² in size and is located between the anterior and posterior meniscofemoral ligaments. The femoral PMB center is located 11.1 mm from the medial arch point and 10.8 mm from the posterior point of the articular cartilage margin. Meanwhile, the PMB tibial attachment center is located 4.4 mm anterior to the champagne glass drop-off of the posterior tibia and 3.1 mm lateral from the medial groove of the medial tibial plateau articular surface (5). These measures have biomechanical and surgical implications, because an anatomic reconstruction of the ALB and PMB better

restores native knee kinematics and has been reported to improve clinical outcomes [Figure 1].

Biomechanics

Functionally, the PCL is a primary restraint to posterior tibial translation at all flexion angles. It also has a role in primary restraint for internal rotation beyond 90° and a supplemental restraint to external tibial rotation beyond 90° of flexion (7). Both bundles have a synergistic and codominant behavior during knee range of motion (ROM) (6, 7). Historically, the ALB and PMB were believed to function independently in a reciprocal nature, with the ALB primarily functioning in deep flexion and the PMB in extension (4, 8, 9). However, recent biomechanical studies have demonstrated that both the ALB and PMB assume a significant role in resisting posterior tibial translation at all flexion angles. This suggests a codominant relationship between both bundles and, therefore, both assume a significant role in knee stability in the absence of the other bundle (4, 6, 10).

The ALB is the main resistant to posterior tibial

translation between 70° and 105°, while the PMB is the main resistant between 0° and 15°. This distribution of forces between the two bundles has surgical implications at the time of graft fixation during anatomic double-bundle (DB) PCL reconstructions (PCLR). Kennedy et al. reported in a biomechanical study that when both bundles were sectioned, 11.7 mm of posterior tibial translation at 90° was observed (7). This suggests that to have a grade III PCL injury, both bundles need to be torn. The PCL has recently been reported to have a more important role for rotational stability than previously thought. It restricts internal rotation at all flexion angles, and the PMB was reported to be the most important bundle for controlling rotation beyond 90° of flexion.

Evaluation

PCL tears are typically produced by external trauma such as the classic “dashboard injury” resulting from a posteriorly directed force on the anterior aspect of the proximal tibia with the knee flexed. In athletics, the typical mechanism of isolated PCL tears is a direct blow to the anterior tibia or a fall onto the knee with the foot in a plantar flexed position. Football, soccer, rugby and skiing are among the sports with highest incidence of PCL tears (1). Non-contact mechanisms, such as hyperflexion or hyperextension, are less common (11). Symptoms depend upon the injury mechanism (high vs low-energy) as well as chronicity. Stiffness, swelling and pain on the posterior aspect of the knee are typical symptoms, while anterior knee pain and instability when descending stairs are more often associated with chronic isolated tears (12).

Physical examination for acute conditions of the knee can be difficult due to pain and guarding. The combination of clinical examination tests, mechanism of injury, and symptoms is vital to make an accurate diagnosis of PCL deficiency. It is critical to examine the contralateral knee first and compare it to the injured knee. The posterior drawer test is performed at 90° of flexion, and has a sensitivity of 90% and a specificity of 99% (5, 13). A false-positive pseudo-Lachman test for the ACL is not uncommon. An important indication of PCL deficiency is a positive Clancy sign, which is a loss of the normal anteromedial and lateral prominences of the tibial plateau beneath the femoral condyles, as determined by palpation with the knee at 90° of flexion and neutral rotation. The posterior sag test (Godfrey test) is performed with the hip and knee flexed to 90° while the examiner supports the leg. If the PCL is torn, an abnormal contour or sag may be evident at the proximal anterior tibia viewed from a lateral position. The quadriceps active test is performed with the patient supine and the knee 90° flexed while the examiner stabilizes the foot. A positive test is observed when the patient performs an isometric quadriceps contraction and dynamically reduces the tibia. The dial test is used to assess combined lesions to the posterolateral corner (PLC) of the knee by assessing external rotation. If it is positive at both 30° and 90° of flexion, a PLC grade III tear combined with a PCL tear

is present (14). In a recent study by Moulton et al, side-to-side differences in internal rotation were assessed under anesthesia by measuring anterior tibial tubercle excursion. The supine IR test performed between 60° and 120° resulted in 95.5% sensitive and 97.1% specific in diagnosing a grade III PCL tear (15).

Additional maneuvers must be utilized to evaluate for possible combined ligament and concomitant intraarticular injury. In a recent systematic review, Kopkow et al, reported that the quadriceps active test is the most specific test for detecting PCL deficiency, and the posterior sag sign is the most sensitive test, although there is a reported high risk of bias among studies reporting diagnosis (16).

Imaging Studies

Standard weightbearing radiographs are performed to detect the presence of fractures, bony avulsions, joint space assessment and tibiofemoral joint congruity. When posterior knee instability is discovered on physical exam, posterior stress radiographs should be performed to objectively quantify posterior knee laxity (17–21). Kneeling stress radiography allows for comparison of the magnitude of posterior tibial displacement on the femur between the injured and uninjured knees. A diagnostic algorithm has been validated where (1) 0–7 mm of side-to-side difference in posterior displacement constitutes a partial PCL tear; (2) 8–11 mm constitutes an isolated complete PCL tear and (3) ≥12 mm of posterior translation constitutes a combined PCL and posterolateral corner or posteromedial corner knee injury [Figure 2] (22).

Magnetic resonance imaging (MRI) is an important adjunct to the diagnosis of PCL tears because it has been found to have a sensitivity, specificity, and accuracy of nearly 100% for the diagnosis of acute PCL injuries (23–25). MRI has lower sensitivity in the evaluation of chronic PCL tears because the signal and shape of the PCL can be deceptively restored through the healing process in chronic cases despite residual laxity being present. However, a recent MRI study by Wilson et al quantified T2 and T2* PCL properties in asymptomatic population (26). They reported significant differences in T2 values in distal, middle and proximal regions of the PCL providing a feasible baseline to compare acute and chronic PCL tears in the future. Therefore, stress radiographs are strongly advocated to diagnose chronic PCL tears (27). MRI is also important to diagnose concurrent meniscal, cartilage, and ligamentous injuries.

In addition, it is important to evaluate alignment with weightbearing long limb radiographs, as well as sagittal plane tibial slope, especially in chronic or revision cases. Patients with isolated PCL tears and a decreased posterior tibial slope may be candidates for a high tibial osteotomy (HTO) to increase their slope and thereby decrease graft forces and reconstruction graft failure rate. Varus and valgus stress radiographs are also helpful in objectively diagnosing suspected concurrent medial and/or lateral sided injuries based on exam findings (28, 29).




Figure 2. Lateral kneeling posterior stress radiographs that demonstrate an increase of 11.6 mm of posterior translation between the injured and uninjured knee. A line is extended parallel from the posterior cortex from at least 15 cm distal to the joint line. A perpendicular line is drawn from this line to the posterior point of the Blumensaat line and the distance is measured and recorded for each knee. The difference between these two points is the posterior tibial translation distance. (*"Jackman T, LaPrade RF, Pontinen T, Lender PA. Intraobserver and interobserver reliability of the kneeling technique of stress radiography for the evaluation of posterior knee laxity. Am J Sports Med. 2008;36:1571-1776."*)

Treatment Rationale

The treatment of complete, isolated PCL tears remains controversial. Some studies have reported good outcomes after conservative treatment of partial PCL tears, while others have reported poor results at long-term follow-up with disabling symptoms and functional limitations (28-35, 30-41). Most authors agree that partial isolated PCL tears should be treated nonoperatively. Complete PCL tears treated nonoperatively have been reported to increase the risk of degenerative changes of the medial and patellofemoral compartments at long term, and were associated with poor function (42-45). Surgical treatment is therefore recommended for symptomatic complete and combined PCL injuries to restore joint stability and improve function. Surgical treatment of PCL tears has also been reported to improve patient outcomes. Several surgical techniques are described in the literature; however, controversy exists on which is superior. Furthermore, no long-term study has been able to demonstrate that PCLR prevents the development of knee OA (46).

Nonoperative treatment

Nonoperative treatment is an option for isolated acute PCL tears because of the inherent healing capacity of the PCL; however, there is a risk that it can heal in a lax position (47-50). Dynamic PCL braces can help keep the tibia reduced during healing by avoiding posterior tibial sag (47, 51, 52). It has been reported that because the PCL has a variable tension throughout knee ROM, a properly designed PCL brace should apply a force that varies with knee flexion angle and replicate anatomic

forces applied to the native PCL. This led to the design of functional dynamic force braces, which provide significantly greater applied force at 45° of flexion that increases with knee flexion angle (51, 52). However, further clinical studies are necessary to determine whether posterior knee laxity is improved long-term following treatment of PCL tears with a dynamic brace. Dynamic bracing is indicated both for nonoperative treatment and postoperative rehabilitation of PCL tears. If nonoperative treatment fails, operative treatment is indicated.

Operative Treatment

Several techniques for PCLR have been described in the literature, depending on tibial graft fixation (transtibial tunnel and tibial inlay techniques), the bundles addressed (single-bundle or double-bundle), and the type of graft used (53-57). The inlay techniques were developed to avoid the sharp angle at the proximal aperture of the tibial tunnel ("killer turn") when using a patellar tendon graft that can damage the PCLR graft, and increase the risk of failure (24, 53, 58). The inlay technique involves creating a trough at the tibial attachment of PCL to match with the graft bone plug fixed with a cannulated screw (with or without washers). Traditional inlay technique requires an open posteromedial approach between the semimembranosus tendon and the medial head of the gastrocnemius muscle, although arthroscopic tibial inlay techniques have been described (53, 58).

One of the biggest controversies concerning PCL




Figure 3. Examination under anesthesia. On the left, a posterior sag is observed. On the right, an anterior drawer is performed to reduce the posterior tibial subluxation.

reconstructions is regarding the outcomes of SB versus DB PCLR. While a SB PCL technique reconstructs only the ALB, a DB PCL technique reconstructs both ALB and PMB, thereby restoring normal anatomy and native knee kinematics (5–7, 27, 54, 57, 59–61). Recent biomechanical studies have demonstrated that a DB PCLR restores knee kinematics to near native better than a single bundle (SB) PCLR. Furthermore, DB PCLR restores rotational stability better than SB PCLR (57).

The authors preferred technique (after an examination under anesthesia), is a DB PCLR with an 11 mm Achilles tendon allograft for the ALB, and a 7 mm tibialis anterior allograft for the PMB [Figure 3]. For femoral tunnels, 11 mm and 7 mm diameter tunnels with a 2 mm bone bridge between the ALB and PMB reconstruction tunnels, respectively, are performed [Figure 4]. On the tibial side, a 12 mm tunnel is reamed under fluoroscopic guidance towards the center of the PCL tibial footprint. It has been demonstrated that aiming solely towards the ALB footprint, especially in SB reconstruction, might injure the medial meniscal root attachment and therefore lead to medial compartment cartilage overload with increased joint contact pressures comparable to a medial meniscectomy (62) [Figure 5]. The ALB is fixed at 90° with an anterior drawer to reduce the normal tibiofemoral step-off, and the PMB is fixed in full extension [Figure 6]. In the chronic setting, a limb alignment assessment is systematically performed to rule out a possible two-stage treatment with a first-

stage corrective osteotomy and a second stage PCLR (63). In a multi-ligament reconstruction injury setting, in order to avoid PCL femoral tunnel convergence, the superficial medial collateral ligament (sMCL) tunnel should be oriented 40° proximally and anteriorly, while the posterior oblique ligament (POL) should be oriented 20° proximally and anteriorly (64). On the tibial side, the POL tunnel should be aimed 15 mm medial to Gerdy's tubercle, while the sMCL tunnel should be oriented transversely across the tibia (anterior to the fibula) and 30° distally in order to avoid tunnel convergence (65).

Post-Op Rehabilitation and Bracing

Following an algorithmic approach to the diagnosis and treatment of a PCL tear, rehabilitation plays a crucial next step in determining patient outcomes (27, 66, 67). Although different rehabilitation programs exist, there are key elements that should lay the foundation for any protocol. These elements include progressive weight-bearing, prevention of posterior tibial subluxation, and early quadriceps strengthening (66–68). We recommend that PCLR patients be kept non-weight bearing for 6 weeks given that PCL graft healing time has been reported to be almost double that following ACL reconstruction (66, 68, 69). Patients are initially placed into a knee immobilizer brace for 3 days prior to transitioning to a dynamic anterior drawer brace. It has been recommended that the PCL brace be worn around the clock for up to a minimum of 24 weeks postoperatively (67). A progressive, goal-oriented, 5-phase rehabilitation program following acute,




Figure 4. (A) Posterior and (B) anterior illustrations of the anatomic double-bundle, posterior cruciate ligament reconstruction. ACL, anterior cruciate ligament; aMFL, anterior meniscomfemoral ligament (ligament of Humphrey); FCL, fibular collateral ligament; PFL, popliteofibular ligament; PMFL, posterior meniscomfemoral ligament (ligament of Wrisberg); POL, posterior oblique ligament (*Reproduced with permission from "Wijdicks CA, Kennedy NI, Goldsmith MT, et al. Kinematic analysis of the posterior cruciate ligament, part 2: a comparison of anatomic single-versus double-bundle reconstruction. Am J Sports Med. 2013;41(12):2839-2948"*).




Figure 5. Fluoroscopic image of transtibial tunnel guide pin placement. On the left, the lateral view shows the guide pin successfully positioned approximately 6 to 7 mm proximal to the champagne-glass drop-off at the PCL facet. On the right, the AP view shows appropriate position of the guidewire at the medial aspect of the lateral tibial eminence and 1 to 2 mm distal to the joint line.




Figure 6. Illustration depicting the intimate relationship of the posterior meniscal roots with the posterior cruciate ligament (PCL) (right knee). LPRA, lateral meniscal posterior root attachment; MPRA, medial meniscal posterior root attachment; SWF, shiny white fibers. Reproduced with permission from "Johannsen AM, Civitarese DM, Padalecki JR, Goldsmith MT, Wijedicks CA, LaPrade RF. Qualitative and quantitative anatomic analysis of the posterior root attachments of the medial and lateral menisci. *Am J Sports Med.* 2012;40(10):2342-2347".

isolated PCLR has been reported to improve stabilization of posterior tibial translation, varus, and external rotation stresses (70). Pierce suggested such a protocol (67). Phase I, 0 to 6 weeks after surgery, is marked by progressive range of motion (ROM) exercises beginning with passive prone ROM from 0 to 90 degrees of knee flexion for the first 2 weeks after surgery advancing to full passive prone ROM as tolerated. During this phase, it is critical to prevent hyperextension and posterior tibial translation to protect the healing PCL graft from elongating. Phase II, from 7 to 12 weeks postoperatively, involves similar precautions with progression to crutch weaning and weightbearing activities as tolerated, while restricting the knee to less than 70° of flexion during weightbearing exercises. Brace use continues in phase III, from 13 to 18 weeks after surgery, with ROM weight-bearing exercise progressing past 70° of knee flexion after 16 weeks. Phase IV, 19 to 24 weeks postoperatively, is characterized by the gradual introduction of sport-specific drills. In phase V, 25 to 36 weeks after surgery, the patient may begin to wean from brace use if the 6 month postoperative PCL stress radiographs demonstrate sufficient healing and begin a straight-line jogging progression with the eventual goal of multiplanar agility exercises and, ultimately, return to preoperative activities (67). Although the above protocol reflects rehab guidelines following an acute isolated PCLR and chronic isolated or combined ligament PCLR may be rehabilitated in a similar fashion, PCL stress radiographs may be required to objectively

gauge postoperative progression and to determine any modifications for a patient with concomitant MCL, PLC, or meniscal injury (22, 27).

Outcomes

Tibial Inlay and Transtibial SB Techniques

A recent systematic review analyzed seven studies between 2006 and 2014 that evaluated the outcome scores for SB tibial inlay and transtibial SB techniques. The authors reported that there were no clinically significant differences in outcomes between both treatments (71). However, 26% of knees in the transtibial group and 27% of knees in the tibial inlay group had Grade II or greater posterior laxity postoperatively. Small discrepancies were found between Lysholm and Tegner scores, but were determined to not be statistically significant in a clinical setting (71). Of the four studies reporting Tegner scores, only one suggested slight superiority for the tibial inlay technique, with the margin being only 0.5 points higher than the transtibial technique (72). For all studies reviewed, Tegner scores ranged from 5.6 to 6 for the transtibial technique and 5.84 to 6.1 for the tibial inlay technique (71). For Lysholm scores, only one of five studies favored the transtibial approach, although this differential was less than 7 points in scale out of 100 (71, 73). Lysholm scores in this systematic review ranged from 81 to 91.3 for the transtibial technique and 76 to 92.8 for the tibial inlay technique (71).

In an earlier systematic review by Kim et al, it was reported that 75% of patients analyzed over 10 studies had normal or nearly normal subjective IKDC scores for those who underwent transtibial PCLR (74). Objectively, posterior knee laxity ranged between 2.0 to 5.9 mm postoperatively among the studies reviewed, which was substantial improvement from preoperative scores ranging between 8.4 mm to 12.3 mm (significance was not reported) (74). Although a large decrease in laxity was seen, it was concluded that normal knee function and posterior stability was not restored in patients analyzed in any of the reviewed studies. Hermans et al. reported that 60% of patients had evidence of osteoarthritis (OA) after SB PCLR at 9.1 years follow up – a possible concomitant occurrence due to the inability to fully restore knee kinematics (54). Therefore, despite improved outcomes after SB procedures, persistent posterior laxity remains a problem, and it has been theorized that this instability can lead to the development of OA.

Clinical Outcomes Comparison between SB and DB PCLR

Recently, Chahla et al performed a systematic review and meta-analysis comparing the SB versus DB PCLR technique. Significantly improved objective posterior tibial stability and objective IKDC scores were obtained within the DB PCLR cohort in comparison to SB group. However, no significant differences were reported in postoperative Lysholm or Tegner scores (75). Another recent systematic review comparing the two techniques challenged the idea that DB techniques were superior to a SB PCLR (76). In their evaluation of eight studies, with levels of evidence (LOE) ranging between II to V, seven studies reported no statistical differences in functional and objective assessments when comparing both techniques (76). Li et al (LOE=2) was the only study suggesting that DB PCLR was superior at a minimum 2-years follow-up. Using a KT-1000 arthrometer, a significant difference was reported in posterior translation of 4.1 and 2.2 mm for SB and DB PCLR, respectively ($P<.05$) (77).

It is worth mentioning that a limitation when comparing these studies falls on the varied utilization of tibial inlay and transtibial surgical procedures. However, recent literature has suggested that the difference between the two techniques is small in the context of a DB PCLR. Four studies utilizing the transtibial technique reported that postoperative subjective outcome substantially improved (19, 65, 69, 70). Additionally, these studies ranged between 0.9 mm to 3.9 mm in posterior translation postoperatively, with the highest values of this range being involved with combined ligamentous injury of the posterolateral corner (74).

Similar to the transtibial technique, studies utilizing the tibial inlay technique reported significant

improvement in subjective and functional outcome scores. In addition, at a minimum follow-up of 2 years, Telos stress radiography showed that posterior translation for isolated PCL injury had a significant improvement of 2.6 to 5.1 mm postoperatively based on two reports (58, 78). In addition, Lee et al recommended that a tibial inlay DB procedure might be the best option for revision PCLR, as it was reported to have the lowest posterior translation of 2.4 mm postoperatively (79). When comparing posterior translation between the two techniques, objective posterior translation measurements via Telos radiograph suggest a slight advantage to the transtibial technique. Therefore, LaPrade et al recommended that the transtibial DB PCLR procedure should be selected given the ability to most closely restore native knee kinematics (27).

Limitations exist when comparing all studies described, as they vary with their assessments of posterior tibial translation by utilizing either a KT-1000 arthrometer, kneeling, or Telos stress radiograph technique. Currently, literature lacks studies with level I of evidence of PCLR. Furthermore, there is a relative paucity between the long-term outcomes of patients who underwent a DB PCLR procedure since this treatment option is still emerging. Future long-term studies should be performed to ultimately distinguish any significant differences between SB and DB surgical treatments and their outcomes.

An improved understanding of PCL tear diagnosis, anatomy, biomechanics, and surgical technique has recently been demonstrated. Stress radiography has become an integral and objective component within the PCL treatment algorithm. Anatomy and biomechanics studies have highlighted the codominant behavior of the ALB and PMB. When restoring both these bundles on the femur with DB PCLR, knee kinematics are also restored. Variability of outcomes measurements between studies have made it difficult to determine clinical differences between SB and DB PCLR; however, recent literature has more strongly substantiated the clinical benefit of DB PCLR over SB technique. Future prospective, long-term outcomes studies are needed as the treatment of PCL tears advances.

Santiago Pache MD
Zachary S. Aman BA
Mitchell Kennedy BA
Gilberto Yoshinobu Nakama MD
Gilbert Moatshe MD
Connor Ziegler MD
Robert F LaPrade PhD
Steadman Philippon Research Institute, The Steadman
Clinic, Vail, Colorado, USA

References

1. Fanelli GC, Edson CJ. Posterior cruciate ligament injuries in trauma patients: Part II. Arthroscopy. 1995; 11(5):526-9.
2. Logan M, Williams A, Lavelle J, Gedroyc W, Freeman M. The effect of posterior cruciate ligament deficiency on knee kinematics. Am J Sports Med. 2004; 32(8):1915-22.
3. MacDonald P, Miniaci A, Fowler P, Marks P, Finlay B. A biomechanical analysis of joint contact forces in the posterior cruciate deficient knee. Knee Surg Sports Traumatol Arthrosc. 1996; 3(4):252-5.
4. Kennedy NI, LaPrade RF, Goldsmith MT, Faucett SC, Rasmussen MT, Coatney GA, et al. Posterior cruciate ligament graft fixation angles, part 1: biomechanical evaluation for anatomic single-bundle reconstruction. Am J Sports Med. 2014; 42(10):2338-45.
5. Anderson CJ, Ziegler CG, Wijdicks CA, Engebretsen L, LaPrade RF. Arthroscopically pertinent anatomy of the anterolateral and posteromedial bundles of the posterior cruciate ligament. J Bone Joint Surg Am. 2012; 94(21):1936-45.
6. Ahmad CS, Cohen ZA, Levine WN, Gardner TR, Ateshian GA, Mow VC. Codominance of the individual posterior cruciate ligament bundles. An analysis of bundle lengths and orientation. Am J Sports Med. 2003; 31(2):221-5.
7. Kennedy NI, Wijdicks CA, Goldsmith MT, Michalski MP, Devitt BM, Årøen A, et al. Kinematic analysis of the posterior cruciate ligament, part 1: the individual and collective function of the anterolateral and posteromedial bundles. Am J Sports Med. 2013; 41(12):2828-38.
8. Boutefnouchet T, Bentayeb M, Qadri Q, Ali S. Long-term outcomes following single-bundle transtibial arthroscopic posterior cruciate ligament reconstruction. Int Orthop. 2013; 37(2):337-43.
9. Dennis MG, Fox JA, Alford JW, Hayden JK, Bach BR Jr. Posterior cruciate ligament reconstruction: current trends. J Knee Surg. 2004; 17(3):133-9.
10. Harner CD, Xerogeanes JW, Livesay GA, Carlin GJ, Smith BA, Kusayama T, et al. The human posterior cruciate ligament complex: an interdisciplinary study. Ligament morphology and biomechanical evaluation. Am J Sports Med. 1995; 23(6):736-45.
11. Strickland JP, Fester EW, Noyes FR. Lateral, posterior, and cruciate knee anatomy. Noyes' knee Disord surgery, Rehabil Clin outcomes. Philadelphia: Saunders; 2009. P. 20-43.
12. Girgis FG, Marshall JL, Monajem A. The cruciate ligaments of the knee joint. Anatomical, functional and experimental analysis. Clin Orthop Relat Res. 1975; 106:216-31.
13. Markolf KL, Feeley BT, Tejwani SG, Martin DE, McAllister DR. Changes in knee laxity and ligament force after sectioning the posteromedial bundle of the posterior cruciate ligament. Arthroscopy. 2006; 22(10):1100-6.
14. Tsai AG, Wijdicks CA, Walsh MP, LaPrade RF. Comparative kinematic evaluation of all-inside single-bundle and double-bundle anterior cruciate ligament reconstruction: a biomechanical study. Am J Sports Med. 2010; 38(2):263-72.
15. Moulton SG, Cram TR, James EW, Dornan GJ, Kennedy NI, LaPrade RF. The Supine internal rotation test: a pilot study evaluating tibial internal rotation in grade III posterior cruciate ligament tears. Orthop J Sport Med. 2015; 3(2):2325967115572135.
16. Kopkow C, Freiberg A, Kirschner S, Seidler A, Schmitt J. Physical examination tests for the diagnosis of posterior cruciate ligament rupture: a systematic review. J Orthop Sports Phys Ther. 2013; 43(11):804-13.
17. Hewett TE, Noyes FR, Lee MD. Diagnosis of complete and partial posterior cruciate ligament ruptures. Stress radiography compared with KT-1000 arthrometer and posterior drawer testing. Am J Sports Med. 1997; 25(5):648-55.
18. Jung TM, Reinhardt C, Scheffler SU, Weiler A. Stress radiography to measure posterior cruciate ligament insufficiency: a comparison of five different techniques. Knee Surg Sports Traumatol Arthrosc. 2006; 14(11):1116-21.
19. Margheritini F, Mariani PP. Diagnostic evaluation of posterior cruciate ligament injuries. Knee Surg Sports Traumatol Arthrosc. 2003; 11(5):282-8.
20. Spiridonov SI, Slinkard NJ, LaPrade RF. Isolated and combined grade-III posterior cruciate ligament tears treated with double-bundle reconstruction with use of endoscopically placed femoral tunnels and grafts: operative technique and clinical outcomes. J Bone Joint Surg Am. 2011; 93(19):1773-80.
21. Stäubli HU, Jakob RP. Posterior instability of the knee near extension. A clinical and stress radiographic analysis of acute injuries of the posterior cruciate ligament. J Bone Joint Surg Br. 1990; 72(2):225-30.
22. Jackman T, LaPrade RF, Pontinen T, Lender PA. Intraobserver and interobserver reliability of the kneeling technique of stress radiography for the evaluation of posterior knee laxity. Am J Sports Med. 2008; 36(8):1571-6.
23. Bedi A, Musahl V, Cowan JB. Management of posterior cruciate ligament injuries: an evidence-based review. J Am Acad Orthop Surg. 2016; 24(5):277-89.
24. Gross ML, Grover JS, Bassett LW, Seeger LL, Finerman GA. Magnetic resonance imaging of the posterior cruciate ligament. Clinical use to improve diagnostic accuracy. Am J Sports Med. 1992; 20(6):732-7.
25. Heron CW, Calvert PT. Three-dimensional gradient-echo MR imaging of the knee: comparison with arthroscopy in 100 patients. Radiology. 1992; 183(3):839-44.
26. Wilson KJ, Surowiec RK, Ho CP, Devitt BM, Fripp J, Smith WS, et al. Quantifiable imaging biomarkers for

- evaluation of the posterior cruciate ligament using 3-T magnetic resonance imaging: a feasibility study. *Orthop J Sport Med.* 2016; 4(4):2325967116639044.
27. LaPrade CM, Civitarese DM, Rasmussen MT, LaPrade RF. Emerging updates on the posterior cruciate ligament: a review of the current literature. *Am J Sports Med.* 2015; 43(12):3077-92.
 28. LaPrade RF, Heikes C, Bakker AJ, Jakobsen RB. The reproducibility and repeatability of varus stress radiographs in the assessment of isolated fibular collateral ligament and grade-III posterolateral knee injuries. An in vitro biomechanical study. *J Bone Joint Surg Am.* 2008; 90(10):2069-76.
 29. LaPrade RF, Bernhardtson AS, Griffith CJ, Macalena JA, Wijdicks CA. Correlation of valgus stress radiographs with medial knee ligament injuries: an in vitro biomechanical study. *Am J Sports Med.* 2010; 38(2):330-8.
 30. Cross MJ, Powell JF. Long-term followup of posterior cruciate ligament rupture: a study of 116 cases. *Am J Sports Med.* 1984; 12(4):292-7.
 31. Fowler PJ, Messieh SS. Isolated posterior cruciate ligament injuries in athletes. *Am J Sports Med.* 1987; 15(6):553-7.
 32. Parolie JM, Bergfeld JA. Long-term results of nonoperative treatment of isolated posterior cruciate ligament injuries in the athlete. *Am J Sports Med.* 1986; 14(1):35-8.
 33. Shelbourne KD, Davis TJ, Patel DV. The natural history of acute, isolated, nonoperatively treated posterior cruciate ligament injuries. A prospective study. *Am J Sports Med.* 1999; 27(3):276-83.
 34. Shelbourne KD, Muthukaruppan Y. Subjective results of nonoperatively treated, acute, isolated posterior cruciate ligament injuries. *Arthroscopy.* 2005; 21(4):457-61.
 35. Shino K, Horibe S, Nakata K, Maeda A, Hamada M, Nakamura N. Conservative treatment of isolated injuries to the posterior cruciate ligament in athletes. *J Bone Joint Surg Br.* 1995; 77(6):895-900.
 36. Tibone JE, Antich TJ, Perry J, Moynes D. Functional analysis of untreated and reconstructed posterior cruciate ligament injuries. *Am J Sports Med.* 1988; 16(3):217-23.
 37. Torg JS, Barton TM, Pavlov H, Stine R. Natural history of the posterior cruciate ligament-deficient knee. *Clin Orthop Relat Res.* 1989; 246:208-16.
 38. Chen CH, Chen WJ, Shih CH. Arthroscopic double-bundled posterior cruciate ligament reconstruction with quadriceps tendon-patellar bone autograft. *Arthroscopy.* 2000; 16(7):780-2.
 39. Dandy DJ, Pusey RJ. The long-term results of unrepaired tears of the posterior cruciate ligament. *J Bone Joint Surg Br.* 1982; 64(1):92-4.
 40. Dejour H, Walch G, Peyrot J, Eberhard P. [The natural history of rupture of the posterior cruciate ligament]. *Rev Chir Orthop Reparatrice Appar Mot.* 1988; 74(1):35-43.
 41. Keller PM, Shelbourne KD, McCarroll JR, Rettig AC. Nonoperatively treated isolated posterior cruciate ligament injuries. *Am J Sports Med.* 1993; 21(1):132-6.
 42. Boynton MD, Tietjens BR. Long-term followup of the untreated isolated posterior cruciate ligament-deficient knee. *Am J Sports Med.* 1996; 24(3):306-10.
 43. Geissler WB, Whipple TL. Intraarticular abnormalities in association with posterior cruciate ligament injuries. *Am J Sports Med.* 1993; 21(6):846-9.
 44. Gill TJ, DeFrate LE, Wang C, Carey CT, Zayontz S, Zarins B, et al. The effect of posterior cruciate ligament reconstruction on patellofemoral contact pressures in the knee joint under simulated muscle loads. *Am J Sports Med.* 2004; 32(1):109-15.
 45. Strobel MJ, Weiler A, Schulz MS, Russe K, Eichhorn HJ. Arthroscopic evaluation of articular cartilage lesions in posterior-cruciate-ligament-deficient knees. *Arthroscopy.* 2003; 19(3):262-8.
 46. Fanelli GC, Giannotti BF, Edson CJ. Arthroscopically assisted combined posterior cruciate ligament/posterior lateral complex reconstruction. *Arthroscopy.* 1996; 12(5):521-30.
 47. Jacobi M, Reischl N, Wahl P, Gautier E, Jakob RP. Acute isolated injury of the posterior cruciate ligament treated by a dynamic anterior drawer brace: a preliminary report. *J Bone Joint Surg Br.* 2010; 92(10):1381-4.
 48. Tewes DP, Fritts HM, Fields RD, Quick DC, Buss DD. Chronically injured posterior cruciate ligament: magnetic resonance imaging. *Clin Orthop Relat Res.* 1997; 335:224-32.
 49. Patel DV, Allen AA, Warren RF, Wickiewicz TL, Simonian PT. The nonoperative treatment of acute, isolated (partial or complete) posterior cruciate ligament-deficient knees: an intermediate-term follow-up study. *HSS J.* 2007; 3(2):137-46.
 50. Shelbourne KD, Clark M, Gray T. Minimum 10-year follow-up of patients after an acute, isolated posterior cruciate ligament injury treated nonoperatively. *Am J Sports Med.* 2013; 41(7):1526-33.
 51. Jansson KS, Costello KE, O'Brien L, Wijdicks CA, LaPrade RF. A historical perspective of PCL bracing. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21(5):1064-70.
 52. LaPrade RF, Smith SD, Wilson KJ, Wijdicks CA. Quantification of functional brace forces for posterior cruciate ligament injuries on the knee joint: an in vivo investigation. *Knee Surg Sports Traumatol Arthrosc.* 2015; 23(10):3070-6.
 53. Berg EE. Posterior cruciate ligament tibial inlay reconstruction. *Arthroscopy.* 1995; 11(1):69-76.
 54. Hermans S, Corten K, Bellemans J. Long-term results of isolated anterolateral bundle reconstructions of the posterior cruciate ligament: a 6- to 12-year follow-up study. *Am J Sports Med.* 2009; 37(8):1499-507.
 55. MacGillivray JD, Stein BE, Park M, Allen AA, Wickiewicz TL, Warren RF. Comparison of tibial inlay versus transtibial techniques for isolated posterior cruciate ligament reconstruction: minimum 2-year follow-up. *Arthroscopy.* 2006; 22(3):320-8.
 56. Panchal HB, Sekiya JK. Open tibial inlay versus arthroscopic transtibial posterior cruciate ligament reconstructions. *Arthroscopy.* 2011; 27(9):1289-95.

57. Wijdicks CA, Kennedy NI, Goldsmith MT, Devitt BM, Michalski MP, Årøen A, et al. Kinematic analysis of the posterior cruciate ligament, part 2: a comparison of anatomic single- versus double-bundle reconstruction. *Am J Sports Med.* 2013; 41(12):2839–48.
58. Bergfeld JA, McAllister DR, Parker RD, Valdevit AD, Kambic HE. A biomechanical comparison of posterior cruciate ligament reconstruction techniques. *Am J Sports Med.* 2001; 29(2):129–36.
59. Kim SJ, Jung M, Moon HK, Kim SG, Chun YM. Anterolateral transtibial posterior cruciate ligament reconstruction combined with anatomical reconstruction of posterolateral corner insufficiency: comparison of single-bundle versus double-bundle posterior cruciate ligament reconstruction over a 2- to 6. *Am J Sports Med.* 2011; 39(3):481–9.
60. Markolf KL, Feeley BT, Jackson SR, McAllister DR. Where should the femoral tunnel of a posterior cruciate ligament reconstruction be placed to best restore anteroposterior laxity and ligament forces? *Am J Sports Med.* 2006; 34(4):604–11.
61. Papannagari R, DeFrate LE, Nha KW, Moses JM, Moussa M, Gill TJ, et al. Function of posterior cruciate ligament bundles during in vivo knee flexion. *Am J Sports Med.* 2007; 35(9):1507–12.
62. LaPrade CM, Smith SD, Rasmussen MT, Hamming MG, Wijdicks CA, Engebretsen L, et al. Consequences of tibial tunnel reaming on the meniscal roots during cruciate ligament reconstruction in a cadaveric model, Part 2: the posterior cruciate ligament. *Am J Sports Med.* 2015; 43(1):207–12.
63. Chahla J, Nitri M, Civitarese D, Dean CS, Moulton SG, LaPrade RF. Anatomic double-bundle posterior cruciate ligament reconstruction. *Arthrosc Tech.* 2016; 5(1):e149–56.
64. Moatshe G, Brady AW, Slette EL, Chahla J, Turnbull TL, Engebretsen L, et al. Multiple ligament reconstruction femoral tunnels: intertunnel relationships and guidelines to avoid convergence. *Am J Sports Med.* 2017; 45(3):563–9.
65. Moatshe G, Slette EL, Engebretsen L, LaPrade RF. Intertunnel relationships in the tibia during reconstruction of multiple knee ligaments: how to avoid tunnel convergence. *Am J Sports Med.* 2016; 44(11):2864–9.
66. Fanelli GC. Posterior cruciate ligament rehabilitation: how slow should we go? *Arthroscopy.* 2008; 24(2):234–5.
67. Pierce CM, O'Brien L, Griffin LW, LaPrade RF. Posterior cruciate ligament tears: functional and postoperative rehabilitation. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21(5):1071–84.
68. Harner CD, Höher J. Evaluation and treatment of posterior cruciate ligament injuries. *Am J Sports Med.* 1998; 26(3):471–82.
69. Bellelli A, Adriani E, Margheritini F, Camillieri G, Della Rocca C, Mariani PP. [Synovial healing in reconstructed cruciate ligaments. Our personal experience compared in single interventions and combined reconstructions]. *Radiol Med.* 1999; 98(6):454–61.
70. Nyland J, Hester P, Caborn DN. Double-bundle posterior cruciate ligament reconstruction with allograft tissue: 2-year postoperative outcomes. *Knee Surg Sports Traumatol Arthrosc.* 2002; 10(5):274–9.
71. Shin YS, Kim HJ, Lee DH. No clinically important difference in knee scores or instability between transtibial and inlay techniques for pcl reconstruction: a systematic review systematic review. *Clin Orthop Relat Res.* 2017; 475(4):1239–48.
72. Seon JK, Song EK. Reconstruction of isolated posterior cruciate ligament injuries: a clinical comparison of the transtibial and tibial inlay techniques. *Arthroscopy.* 2006; 22(1):27–32.
73. Kim SJ, Kim TE, Jo SB, Kung YP. Comparison of the clinical results of three posterior cruciate ligament reconstruction techniques. *J Bone Joint Surg Am.* 2009; 91(11):2543–9.
74. Kim YM, Lee CA, Matava MJ. Clinical results of arthroscopic single-bundle transtibial posterior cruciate ligament reconstruction: a systematic review. *Am J Sports Med.* 2011; 39(2):425–34.
75. Chahla J, Moatshe G, Cinque ME, Dornan GJ, Mitchell JJ, Ridley TJ, et al. Single bundle and double bundle posterior cruciate ligament reconstructions: a systematic review and meta-analysis of 441 patients at a minimum 2 years follow up. *Arthrosc.* 2017; 33(11):2066–80.
76. Qi YS, Wang HJ, Wang SJ, Zhang ZZ, Huang AB, Yu JK. A systematic review of double-bundle versus single-bundle posterior cruciate ligament reconstruction. *BMC Musculoskelet Disord.* 2016; 17:45–54.
77. Li Y, Li J, Wang J, Gao S, Zhang Y. Comparison of single-bundle and double-bundle isolated posterior cruciate ligament reconstruction with allograft: a prospective, randomized study. *Arthroscopy.* 2014; 30(6):695–700.
78. Noyes FR, Barber-Westin S. Posterior cruciate ligament replacement with a two-strand quadriceps tendon-patellar bone autograft and a tibial inlay technique. *J Bone Joint Surg Am.* 2005; 87(6):1241–52.
79. Lee SH, Jung YB, Lee HJ, Jung HJ, Kim SH. Revision posterior cruciate ligament reconstruction using a modified tibial-inlay double-bundle technique. *J Bone Joint Surg Am.* 2012; 94(6):516–22.